On a differential operator with absent spectrum

Alexander Makin

Abstract

In this paper, we consider spectral problem for the nth order ordinary differential operator with degenerate boundary conditions. We construct a nontrivial example of boundary value problem which has no eigenvalues.

arXiv:1703.09642v1 [math.SP] 15 Mar 2017

Consider the boundary value problem for the nth order ordinary differential equation

$$l(u) = u^{(n)}(x) + \sum_{m=1}^{n} p_m(x) u^{(n-m)}(x) = \lambda u(x) \tag{1}$$

with boundary conditions

$$B_j(u) = u^{(n-j)}(0) + d(-1)^{j+1} u^{(n-j)}(1) = 0 \tag{2}$$

$(j = 1, \dots n)$, where $n = 2\nu$, coefficient d is an arbitrary complex number, and the complex-valued coefficients $p_m(x)$ are functions in the class $L_1(0,1)$. Suppose that $p_m(x) = (-1)^m p_m(1-x)$ almost everywhere on the segment $[0,1]$, $m = 1, \dots, n$. We will study the spectrum of problem (1), (2).

Let a function $u_k(x)$ be the solution of the Cauchy problem

$$l(u) = \lambda u, \quad u_k^{(j)}(1/2) = \delta_{k,j}, \tag{3}$$

where $k = 0, \dots, n-1$, $j = 0, \dots, n-1$. Denote

$$u_-(x) = \sum_{k=0}^{\nu-1} c_{2k+1} u_{2k+1}(x), \quad u_+(x) = \sum_{k=0}^{\nu-2} c_{2k} u_{2k}(x),$$

where c_i are arbitrary constants $(i = 0, \dots, n-1)$. Then

$$u_-^{(2k)}(1/2) = 0, \quad u_+^{(2k+1)}(1/2) = 0, \quad k = 0, \dots, \nu - 1.$$

2000 Mathematics Subject Classification. 34L20.
Key words and phrases. Ordinary differential operator, boundary value problem, spectrum.

Obviously, that the functions $w_-(x) = -u_-(1-x)$ and $w_+(x) = u_+(1-x)$ are the solutions of equation (1) and satisfy the same initial conditions at the point $1/2$ just as the functions $u_-(x)$ and $u_+(x)$, correspondingly. This, together with the uniqueness of the solution of Cauchy problem (3) implies, that $u_-(x) = -u_-(1-x), \quad u_+(x) = u_+(1-x)$ for $0 \le x \le 1$. It follows from this that

$$u_-^{(n-j)}(0) + (-1)^{j+1} u_-^{(n-j)}(1) = 0, \quad u_+^{(n-j)}(0) + (-1)^j u_+^{(n-j)}(1) = 0 \quad (4)$$

$(j = 1, \dots, n)$. It follows from (4) that for any complex number λ the function $u_-(x)$ is a solution of problem (1), (2) if $d = 1$, and for any complex number λ the function $u_+(x)$ is a solution of problem (1), (2) if $d = -1$. Thus, we establish that if $d = \pm 1$ the spectrum of problem (1), (2) fills all complex plane. If $p_m(x) \in C^{n-m}(0,1)$, $m = 1, \dots, n$, this assertion was proved in [1].

Assume, for a number λ a function $\tilde{u}(x)$ is a solution of problem (1), (2) if $d \ne \pm 1$. Then $\tilde{u}(x) = u_+(x) + u_-(x)$. We see that

$$u_-^{(n-j)}(0) + u_+^{(n-j)}(0) + (-1)^{j+1} d (u_-^{(n-j)}(1) + u_+^{(n-j)}(1)) = 0 \quad (5)$$

$(j = 1, \dots, n)$. It follows from (4), (5) that

$$u_-^{(n-j)}(0)(1-d) + u_+^{(n-j)}(0)(1+d) = 0$$

$(j = 1, \dots, n)$. From this and the definition of the functions $u_+(x)$ and $u_-(x)$, we have

$$(1+d) \sum_{k=0}^{\nu-2} c_{2k} u_{2k}^{(n-j)}(0) + (1-d) \sum_{k=0}^{\nu-1} c_{2k+1} u_{2k+1}^{(n-j)}(0) = 0$$

$(j = 1, \dots, n)$, hence, the constants c_i $(i = 0, \dots, n-1)$ satisfy the system of linear equations

$$\sum_{i=0}^{n-1} c_i (1 + d(-1)^i) u_i^{(n-j)}(0) = 0 \quad (6)$$

$(j = 1, \ldots, n)$. The determinant of linear system (6) is

$$\Delta = (1 - d^2)^\nu \det ||u_i^{(n-j)}(0)||.$$

Since the last determinant is the Wronskian of the fundamental system of the solutions of equation (1), it is nonzero. Therefore, system (6) has only trivial solution, i.e. the function $\tilde{u}(x) \equiv 0$. Hence, if $d \neq \pm 1$ problem (1), (2) has no eigenvalues.

For the first time problem (1), (2) for $n = 2$ was investigated in [2].

References

1. V.A. Sadovnichy, B.E. Kanguzhin. On a connection between the spectrum of a differential operator with symmetric coefficients and boundary conditions. Dokl. Akad. Nauk SSSR. 1982. V. 267. No. 2. P. 310-313.

2. M.H. Stone. Irregular differential systems of order two and the related expansion problems. Trans. Amer. Math. Soc. 1927. V. 29. No. 1. P. 23-53.

E-mail: alexmakin@yandex.ru